

TITLE OF THE INVENTION

METHOD FOR PRODUCING L-GLUTAMIC ACID BY FERMENTATION

ACCOMPANIED BY PRECIPITATION

5 BACKGROUND OF THE INVENTION

The present invention relates to a method for producing L-glutamic acid by fermentation accompanied by precipitation. L-Glutamic acid is widely used as a material of seasonings and so forth.

- 10 L-Glutamic acid is mainly produced by fermentative methods using so-called coryneform bacteria producing L-glutamic acid and belonging to the genus *Brevibacterium*, *Corynebacterium* or *Microbacterium*, or mutant strains thereof (Amino Acid Fermentation, pp.195-215, Gakkai Shuppan Center, 15 1986). As methods for producing L-glutamic acid by fermentation by using other bacterial strains, there are known a method using a microorganism belonging to the genus *Bacillus*, *Streptomyces*, *Penicillium* or the like (U.S. Patent No. 3,220,929), a method using a microorganism belonging to 20 the genus *Pseudomonas*, *Arthrobacter*, *Serratia*, *Candida* or the like (U.S. Patent No. 3,563,857), a method using a microorganism belonging to the genus *Bacillus*, *Pseudomonas*, *Serratia*, *Aerobacter aerogenes* (currently referred to as *Enterobacter aerogenes*) or the like (Japanese Patent 25 Publication (Kokoku) No. 32-9393), a method using a mutant strain of *Escherichia coli* (Japanese Patent Application Laid-open (Kokai) No. 5-244970) and so forth. In addition, the inventors of the present invention have proposed a method

for producing L-glutamic acid by using a microorganism belonging to the genus *Klebsiella*, *Erwinia* or *Pantoea* (Japanese Patent Application Laid-open No. 2000-106869).

Further, there have been disclosed various techniques for improving L-glutamic acid-producing ability by enhancing activities of L-glutamic acid biosynthetic enzymes through use of recombinant DNA techniques. For example, it has been reported that introduction of a gene coding for citrate synthase derived from *Escherichia coli* or *Corynebacterium glutamicum* was effective for enhancement of L-glutamic acid-producing ability in *Corynebacterium* or *Brevibacterium* bacteria (Japanese Patent Publication No. 7-121228). In addition, Japanese Patent Application Laid-open No. 61-268185 discloses a cell harboring recombinant DNA containing a glutamate dehydrogenase gene derived from *Corynebacterium* bacteria. Further, Japanese Patent Application Laid-open No. 63-214189 discloses a technique for improving L-glutamic acid-producing ability by amplifying a glutamate dehydrogenase gene, an isocitrate dehydrogenase gene, an aconitate hydratase gene and a citrate synthase gene.

Although L-glutamic acid productivity has been considerably increased by breeding of the aforementioned microorganisms or improvement of production methods, development of methods for more efficiently producing L-glutamic acid at a lower cost is required to respond to further increase of the demand in future.

There is known a method wherein fermentation is performed with crystallizing L-amino acid accumulated in

culture (Japanese Patent Application Laid-open No. 62-288). In this method, the L-amino acid concentration in the culture is maintained below a certain level by precipitating the accumulated L-amino acid in the culture. Specifically,

5 L-tryptophan, L-tyrosine or L-leucine is precipitated during fermentation by adjusting temperature and pH of the culture or adding a surface active agent to the medium.

While a fermentative method with precipitating L-amino acid is known as described above, amino acids suitable for

10 this method are those of relatively low water solubility, and no example of applying the method to highly water-soluble amino acids such as L-glutamic acid is known. In addition, the medium must have low pH to precipitate L-glutamic acid. However, L-glutamic acid-producing bacteria such as those

15 mentioned above cannot grow under acidic conditions, and therefore L-glutamic acid fermentation is performed under neutral conditions (U.S. Patent Nos. 3,220,929 and 3,032,474; Chao K.C. & Foster J.W., J. Bacteriol., 77, pp.715-725 (1959)). Thus, production of L-glutamic acid by

20 fermentation accompanied by precipitation is not known. Furthermore, it is known that growth of most acidophile bacteria is inhibited by organic acids such as acetic acid, lactic acid and succinic acid (Yasuro Oshima Ed., "Extreme Environment Microorganism Handbook", p.231, Science Forum;

25 Borichewski R.M., J. Bacteriol., 93, pp.597-599 (1967) etc.). Therefore, it is considered that many microorganisms are susceptible to L-glutamic acid, which is also an organic acid, under acidic conditions, and there has been no report that

search of microorganisms showing L-glutamic acid-producing ability under acidic conditions was attempted.

SUMMARY OF THE INVENTION

5 In the aforementioned current situation, an object of the present invention is to search and breed a microorganism that produces L-glutamic acid under low pH conditions and to provide a method for producing L-glutamic acid using an obtained microorganism by fermentation with precipitating
10 L-glutamic acid.

 The inventors of the present invention considered during the study for improvement of L-glutamic acid productivity by fermentation that inhibition of the production by L-glutamic acid accumulated in a medium at a
15 high concentration was one of obstructions to the improvement of productivity. For example, cells have an excretory system and an uptake system for L-glutamic acid. However, if L-glutamic acid once excreted into the medium is incorporated into cells again, not only the production
20 efficiency falls, but also the L-glutamic acid biosynthetic reactions are inhibited as a result. In order to avoid the inhibition of production by such accumulation of L-glutamic acid at high concentration, the inventors of the present invention screened microorganisms that can proliferate under
25 acidic conditions and in the presence of a high concentration of L-glutamic acid. As a result, they successfully isolated microorganisms having such properties from a soil, and thus accomplished the present invention.

Thus, the present invention provides the followings.

(1) A microorganism which can metabolize a carbon source at a specific pH in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source, and has ability to accumulate L-glutamic acid in an amount exceeding the amount corresponding to the saturation concentration in the liquid medium at the pH.

(2) The microorganism according to (1), which can grow in the liquid medium.

(3) The microorganism according to (1) or (2), wherein the pH is not more than 5.0.

(4) The microorganism according to any one of (1) to (3), which has at least one of the following characteristics:

(a) the microorganism is enhanced in activity of an enzyme that catalyzes a reaction for biosynthesis of L-glutamic acid; and

(b) the microorganism is decreased in or deficient in activity of an enzyme that catalyzes a reaction branching from a biosynthetic pathway of L-glutamic acid and producing a compound other than L-glutamic acid.

(5) The microorganism according to (4), wherein the enzyme that catalyzes the reaction for biosynthesis of L-glutamic acid is at least one selected from citrate synthase, phosphoenolpyruvate carboxylase and glutamate dehydrogenase.

(6) The microorganism according to (4) or (5), wherein the enzyme that catalyzes the reaction branching from the biosynthetic pathway of L-glutamic acid and producing a

compound other than L-glutamic acid is α -ketoglutarate dehydrogenase.

(7) The microorganism according to any one of (1) to (6), wherein the microorganism belongs to the genus *Enterobacter*.

5 (8) The microorganism according to (7), which is *Enterobacter agglomerans*.

(9) The microorganism according to (8), which has a mutation that causes less extracellular secretion of a viscous material compared with a wild strain when cultured
10 in a medium containing a saccharide.

(10) A method for producing L-glutamic acid by fermentation, which comprises culturing a microorganism as defined in any one of (1) to (9) in a liquid medium of which pH is adjusted to a pH at which L-glutamic acid is precipitated, to produce
15 and accumulate L-glutamic acid and precipitate L-glutamic acid in the medium.

(11) A method for screening a microorganism suitable for producing L-glutamic acid by fermentation with precipitating L-glutamic acid in a liquid medium, which comprises
20 inoculating a sample containing microorganisms into an acidic medium containing L-glutamic acid at a saturation concentration and a carbon source, and selecting a strain that can metabolize the carbon source.

(12) The method according to (11), wherein a strain that
25 can grow in the medium is selected as the strain that can metabolize the carbon source.

(13) The method according to (11) or (12), wherein a pH of the medium is not more than 5.0.

B
B

B
B

SUB
D'

B SUB
D2

20 from *Enterobacter agglomerans* and that derived from
Escherichia coli.

Fig. 4 shows comparison of the amino acid sequence deduced from the nucleotide sequence of the *sdhB* gene derived from *Enterobacter agglomerans* (SEQ ID NO: 10) and that derived from *Escherichia coli* (SEQ ID NO: 11).

Fig. 5 shows comparison of the amino acid sequence deduced from the nucleotide sequence of the *sucC* gene derived from *Enterobacter agglomerans* (SEA ID NO. 2) and that derived from

8
(SEQUENCE NO. 12)
Escherichia coli.

Fig. 6 shows construction of plasmid pMWCPG having a *gltA* gene, a *ppc* gene and a *gdhA* gene.

Fig. 7 shows construction of plasmid RSF-Tet having the replication origin of the broad host spectrum plasmid RSF1010 and a tetracycline resistance gene.

Fig. 8 shows construction of plasmid RSFCPG having the replication origin of the broad host spectrum plasmid RSF1010, a tetracycline resistance gene, a *gltA* gene, a *ppc* gene and a *gdhA* gene.

Fig. 9 shows construction of plasmid pSTVCB having a *gltA* gene.

DETAILED DESCRIPTION OF THE INVENTION

Hereafter, the present invention will be explained in detail.

The microorganism of the present invention is a microorganism that (1) can metabolize a carbon source at a specific pH in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source and (2) has ability to accumulate L-glutamic acid in an amount exceeding the amount corresponding to the saturation concentration in the liquid medium at the pH.

The term "saturation concentration" means a concentration of L-glutamic acid dissolved in a liquid medium when the liquid medium is saturated with L-glutamic acid.

Hereafter, a method for screening a microorganism that can metabolize a carbon source in a liquid medium containing

L-glutamic acid at a saturation concentration and the carbon source at a specific pH will be described. A sample containing microorganisms is inoculated into a liquid medium containing L-glutamic acid at a saturation concentration and a carbon source at a specific pH, and a strain that can metabolize the carbon source is selected. The specific pH is not particularly limited, but is usually not more than about 5.0, preferably not more than about 4.5, more preferably not more than about 4.3. The microorganism of the present invention is used to produce L-glutamic acid by fermentation with precipitating L-glutamic acid. If the pH is too high, it becomes difficult to allow the microorganism to produce L-glutamic acid enough for precipitation. Therefore, pH is preferably in the aforementioned range.

If pH of an aqueous solution containing L-glutamic acid is lowered, the solubility of L-glutamic acid significantly falls around pKa of γ -carboxyl group (4.25, 25°C). The solubility becomes the lowest at the isoelectric point (pH 3.2) and L-glutamic acid exceeding the amount corresponding to the saturation concentration is precipitated. While it depends on the medium composition, L-glutamic acid is usually dissolved in an amount of 10 to 20 g/L at pH 3.2, 30 to 40 g/L at pH 4.0 and 50 to 60 g/L at pH 4.7, at about 30°C. Usually pH does not need to be made below 3.0, because the L-glutamic acid precipitating effect plateaus when pH goes below a certain value. However, pH may be below 3.0.

In addition, the expression that a microorganism "can metabolize the carbon source" means that it can proliferate

or can consume the carbon source even though it cannot proliferate, that is, it indicates that it catabolizes carbon sources such as saccharides or organic acids. Specifically, for example, if a microorganism proliferates when cultured
5 in a liquid medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, still more preferably pH 4.0 at an appropriate temperature, for example, 28°C, 37°C or 50°C for 2 to 4 days, this microorganism can metabolize the
10 carbon source in the medium. Further, for example, even if a microorganism does not proliferate when it is cultured in a liquid medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, still more preferably pH 4.0
15 at an appropriate temperature, for example, 28°C, 37°C or 50°C for 2 to 4 days, the microorganism which consumes the carbon source in the medium is that can metabolize the carbon source in the medium.

The microorganism which can metabolize the carbon
20 source includes a microorganism which can grow in the liquid medium.

The expression that a microorganism "can grow" means that it can proliferate or can produce L-glutamic acid even though it cannot proliferate. Specifically, for example,
25 if a microorganism proliferates when cultured in a liquid medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, still more preferably pH 4.0

at an appropriate temperature, for example, 28°C, 37°C or 50°C for 2 to 4 days, this microorganism can grow in the medium. Further, for example, even if a microorganism does not proliferate when it is cultured in a liquid synthetic medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, still more preferably pH 4.0 at an appropriate temperature, for example, 28°C, 37°C or 50°C for 2 to 4 days, the microorganism which increases the amount of L-glutamic acid in the medium is that can grow in the medium.

The selection described above may be repeated two or more times under the same conditions or with changing pH or the concentration of L-glutamic acid. An initial selection can be performed in a medium containing L-glutamic acid at a concentration lower than the saturation concentration, and thereafter a subsequent selection can be performed in a medium containing L-glutamic acid at a saturation concentration. Further, strains with favorable properties such as superior proliferation rate may be selected.

In addition to the property described above, the microorganism of the present invention has ability to accumulate L-glutamic acid in an amount exceeding the amount corresponding to the saturation concentration of L-glutamic acid in a liquid medium. The pH of the aforementioned liquid medium is preferably the same as or close to that of the medium used for screening a microorganism having the aforementioned property (1). Usually, a microorganism becomes susceptible to L-glutamic acid at a high concentration as pH becomes lower.

Therefore, it is preferred that pH is not low from the viewpoint of resistance to L-glutamic acid, but low pH is preferred from the viewpoint of production of L-glutamic acid with precipitating it. To satisfy these conditions, pH may
5 be in the range of 3 to 5, preferably 4 to 5, more preferably 4.0 to 4.7, still more preferably 4.0 to 4.5, particularly preferably 4.0 to 4.3.

As the microorganism of the present invention or breeding materials therefor, there can be mentioned, for
10 example, microorganisms belonging to the genus *Enterobacter*, *Klebsiella*, *Serratia*, *Pantoea*, *Erwinia*, *Escherichia*, *Corynebacterium*, *Alicyclobacillus*, *Bacillus*, *Saccharomyces* or the like. Among these, microorganisms belonging to the genus *Enterobacter* are preferred. Hereafter, the
15 microorganism of the present invention will be explained mainly for microorganisms belonging to the genus *Enterobacter*, but the present invention can be applied to microorganism belonging to other genera and not limited to the genus *Enterobacter*.

20 As microorganisms belonging to the *Enterobacter*, there can be specifically mentioned *Enterobacter agglomerans*, preferably the *Enterobacter agglomerans* AJ13355 strain. This strain was isolated from a soil in Iwata-shi, Shizuoka, Japan as a strain that can proliferate in a medium containing
25 L-glutamic acid and a carbon source at low pH.

The physiological properties of AJ13355 are as follows:

(1) Gram staining: negative

- (2) Behavior against oxygen: facultative anaerobic
- (3) Catalase: positive
- (4) Oxidase: negative
- (5) Nitrate-reducing ability: negative
- 5 (6) Voges-Proskauer test: positive
- (7) Methyl Red test: negative
- (8) Urease: negative
- (9) Indole production: positive
- (10) Motility: motile
- 10 (11) H₂S production in TSI medium: weakly active
- (12) β -galactosidase: positive
- (13) Saccharide-assimilating property:
 - Arabinose: positive
 - Sucrose: positive
 - 15 Lactose: positive
 - Xylose: positive
 - Sorbitol: positive
 - Inositol: positive
 - Trehalose: positive
 - 20 Maltose: positive
 - Glucose: positive
 - Adonitol: negative
 - Raffinose: positive
 - Salicin: negative
 - 25 Melibiose: positive
- (14) Glycerol-assimilating property: positive
- (15) Organic acid-assimilating property:
 - Citric acid: positive

Tartaric acid: negative

Gluconic acid: positive

Acetic acid: positive

Malonic acid: negative

- 5 (16) Arginine dehydratase: negative
 (17) Ornithine decarboxylase: negative
 (18) Lysine decarboxylase: negative
 (19) Phenylalanine deaminase: negative
 (20) Pigment formation: yellow

- 10 (21) Gelatin liquefaction ability: positive
 (22) Growth pH: growth is possible at pH 4.0, good
 growth at pH 4.5 to 7
 (23) Growth temperature: good growth at 25°C, good
 growth at 30°C, good growth at 37°C, growth is possible at
15 42°C, growth is not possible at 45°C

Based on these bacteriological properties, AJ13355 was determined as *Enterobacter agglomerans*.

- The *Enterobacter agglomerans* AJ13355 was deposited at
20 the National Institute of Bioscience and Human-Technology,
 Agency of Industrial Science and Technology, Ministry of
 International Trade and Industry (postal code: 305-8566, 1-3,
 Higashi 1-chome, Tsukuba-shi, Ibaraki, Japan) on February
 19, 1998 and received an accession number of FERM P-16644.
25 It was then transferred to an international deposition under
 the provisions of Budapest Treaty on January 11, 1999 and
 received an accession number of FERM BP-6614.

The microorganism of the present invention may be a

microorganism originally having L-glutamic acid-producing ability or one having L-glutamic acid-producing ability imparted or enhanced by breeding through use of mutation treatment, recombinant DNA techniques or the like.

- 5 L-Glutamic acid-producing ability can be imparted or enhanced by, for example, increasing activity of an enzyme that catalyzes a reaction for biosynthesis of L-glutamic acid. L-Glutamic acid-producing ability can also be enhanced by decreasing activity of an enzyme that catalyzes a reaction
10 branching from the biosynthetic pathway of L-glutamic acid and producing a compound other than L-glutamic acid, or making the activity deficient.

- As enzymes that catalyze a reaction for biosynthesis of L-glutamic acid, there can be mentioned glutamate
15 dehydrogenase (hereafter, also referred to as "GDH"), glutamine synthetase, glutamate synthase, isocitrate dehydrogenase, aconitate hydratase, citrate synthase (hereafter, also referred to as "CS"), phosphoenolpyruvate carboxylase (hereafter, also referred to as "PEPC"),
20 pyruvate dehydrogenase, pyruvate kinase, enolase, phosphoglyceromutase, phosphoglycerate kinase, glyceraldehyde-3-phosphate dehydrogenase, triosephosphate isomerase, fructose bisphosphate aldolase, phosphofructokinase, glucose phosphate isomerase and so
25 forth. Among these enzymes, one, two or three of CS, PEPC and GDH are preferred. Further, it is preferred that the activities of all the three enzymes, CS, PEPC and GDH, are enhanced in the microorganism of the present invention. In

particular, CS of *Brevibacterium lactofermentum* is preferred, because it does not suffer from inhibition by α -ketoglutaric acid, L-glutamic acid and NADH.

In order to enhance the activity of CS, PEPC or GDH, for example, a gene coding for CS, PEPC or GDH may be cloned on an appropriate plasmid and a host microorganism may be transformed with the obtained plasmid. The copy number of the gene coding for CS, PEPC or GDH (hereafter, abbreviated as "*gltA* gene", "*ppc* gene" and "*gdhA* gene", respectively) in the transformed strain cell increases, resulting in the increase of the activity of CS, PEPC or GDH.

The cloned *gltA* gene, *ppc* gene and *gdhA* gene are introduced into the aforementioned starting parent strain solely or in combination of arbitrary two or three kinds of them. When two or three kinds of the genes are introduced, two or three kinds of the genes may be cloned on one kind of plasmid and introduced into the host, or separately cloned on two or three kinds of plasmids that can coexist and introduced into the host.

Two or more kinds of genes coding for enzymes of the same kind, but derived from different microorganisms may be introduced into the same host.

The plasmids described above are not particularly limited so long as they are autonomously replicable in cells of a microorganism belonging to, for example, the genus *Enterobacter* or the like, but, for example, there can be mentioned pUC19, pUC18, pBR322, pHSG299, pHSG298, pHSG399, pH SG398, RSF1010, pMW119, pMW118, pMW219, pMW218, pACYC177,

pACYC184 and so forth. Besides these, vectors of phage DNA can also be used.

Transformation can be performed by, for example, the method of D.M. Morrison (Methods in Enzymology, 68, 326 (1979)), the method wherein permeability of DNA is increased by treating recipient bacterium cells with calcium chloride (Mandel M. and Higa A., J. Mol. Biol., 53, 159 (1970)), the electroporation (Miller J.H., "A Short Course in Bacterial Genetics", Cold Spring Harbor Laboratory Press, U.S.A. 1992) or the like.

The activity of CS, PEPC or GDH can also be increased by allowing multiple copies of a *gltA* gene, a *ppc* gene or a *gdhA* gene to be present on chromosomal DNA of the aforementioned starting parent strain to be a host. In order to introduce multiple copies of the *gltA* gene, the *ppc* gene or the *gdhA* gene on chromosomal DNA of a microorganism belonging to the genus *Enterobacter* or the like, a sequence of which multiple copies are present on the chromosomal DNA, such as repetitive DNA and inverted repeats present at termini of a transposable element, can be used.

Alternatively, multiple copies of the genes can be introduced onto chromosomal DNA by utilizing transfer of a transposon containing the *gltA* gene, the *ppc* gene or the *gdhA* gene. As a result, the copy number of the *gltA* gene, the *ppc* gene or the *gdhA* gene in a transformed strain cell is increased, and thus the activity of CS, PEPC or GDH is increased.

As organisms to be a source of the *gltA* gene, the *ppc* gene or the *gdhA* gene of which copy number is increased, any

organism can be used so long as it has activity of CS, PEPC or GDH. Inter alia, bacteria, which are prokaryotes, for example, those belonging to the genus *Enterobacter*, *Klebsiella*, *Erwinia*, *Pantoea*, *Serratia*, *Escherichia*,

- 5 *Corynebacterium*, *Brevibacterium* and *Bacillus* are preferred. As specific examples, there can be mentioned *Escherichia coli*, *Brevibacterium lactofermentum* and so forth. The *gltA* gene, the *ppc* gene and the *gdhA* gene can be obtained from chromosomal DNA of the microorganisms described above.

- 10 The *gltA* gene, the *ppc* gene and the *gdhA* gene can be obtained by using a mutant strain which is deficient in the activity of CS, PEPC or GDH to isolate a DNA fragment which complements the auxotrophy from chromosomal DNA of the
15 aforementioned microorganisms. Since the nucleotide sequences of these genes of *Escherichia* and *Corynebacterium* bacteria have already been elucidated (Biochemistry, 22, pp.5243-5249 (1983); J. Biochem., 95, pp.909-916 (1984); Gene, 27, pp.193-199 (1984); Microbiology, 140, pp.1817-1828 (1994); Mol. Gen. Genet., 218, pp.330-339 (1989);
20 Molecular Microbiology, 6, pp.317-326 (1992)), they can also be obtained by PCR utilizing primers synthesized based on each nucleotide sequence and chromosomal DNA as a template.

- The activity of CS, PEPC or GDH can also be increased by enhancing the expression of the *gltA* gene, the *ppc* gene
25 or the *gdhA* gene besides the aforementioned amplification of the genes. For example, the expression can be enhanced by replacing a promoter for the *gltA* gene, the *ppc* gene or the *gdhA* gene with other stronger promoters. For example,

lac promoter, *trp* promoter, *trc* promoter, *tac* promoter, P_R promoter and P_L promoter of the lamda phage and so forth are known as strong promoters. The *gltA* gene, the *ppc* gene and the *gdhA* gene of which promoter is replaced are cloned on
5 a plasmid and introduced into the host microorganism, or introduced onto the chromosomal DNA of the host microorganism by using repetitive DNA, inverted repeats, transposon or the like.

The activity of CS, PEPC or GDH can also be enhanced
10 by replacing the promoter of the *gltA* gene, the *ppc* gene or the *gdhA* gene on the chromosome with other stronger promoters (see WO 87/03006 and Japanese Patent Application Laid-open No. 61-268183), or inserting a strong promoter in the upstream of the coding sequence of each gene (see Gene, 29,
15 pp.231-241 (1984)). Specifically, homologous recombination can be performed between DNA containing the *gltA* gene, the *ppc* gene or the *gdhA* gene of which promoter is replaced with a stronger one or a part thereof and the corresponding gene on the chromosome.

20 Examples of the enzyme which catalyze a reaction branching from the biosynthetic pathway of the L-glutamic acid and producing a compound other than L-glutamic acid include α -ketoglutarate dehydrogenase (hereafter, also referred to as " α KGDH"), isocitrate lyase, phosphate
25 acetyltransferase, acetate kinase, acetohydroxy acid synthase, acetolactate synthase, formate acetyltransferase, lactate dehydrogenase, glutamate decarboxylase, 1-pyrroline dehydrogenase and so forth. Among these enzymes,

α KGDH is preferred.

In order to obtain decrease or deficiency of the activity of the aforementioned enzyme in a microorganism belonging to the genus *Enterobacter* or the like, mutation
5 causing decrease or deficiency of the intracellular activity of the enzyme can be introduced into the gene of the aforementioned enzyme by a usual mutagenesis or genetic engineering method.

Examples of the mutagenesis method include, for
10 example, methods utilizing irradiation with X-ray or ultraviolet ray, methods utilizing treatment with a mutagenic agent such as N-methyl-N'-nitro-N-nitrosoguanidine, and so forth. The site where the mutation is introduced to the gene may be in a coding region coding
15 for an enzyme protein, or a region for regulating expression such as a promoter.

Examples of the genetic engineering methods include, for example, methods utilizing gene recombination, transduction, cell fusion and so forth. For example, a drug
20 resistance gene is inserted into a cloned target gene to prepare a gene that has lost its function (defective gene). Subsequently, this defective gene is introduced into a cell of a host microorganism, and the target gene on the chromosome is replaced with the aforementioned defective gene by
25 utilizing homologous recombination (gene disruption).

Decrease or deficiency of intracellular activity of the target enzyme and the degree of decrease of the activity can be determined by measuring the enzyme activity of a cell

extract or a purified fraction thereof obtained from a candidate strain and comparing with that of a wild strain. For example, the α KGDH activity can be measured by the method of Reed et al. (Reed L.J. and Mukherjee B.B., Methods in Enzymology, 13, pp.55-61 (1969)).

Depending on the target enzyme, the target mutant strain can be selected based on the phenotype of the mutant strain. For example, a mutant strain which is deficient in the α KGDH activity or decreases in the α KGDH activity cannot proliferate or shows a markedly reduced proliferation rate in a minimal medium containing glucose or a minimal medium containing acetic acid or L-glutamic acid as an exclusive carbon source under aerobic conditions. However, normal proliferation is enabled even under the same condition by adding succinic acid or lysine, methionine and diaminopimelic acid to a minimal medium containing glucose. By utilizing these phenomena as indicators, mutant strains with decreased α KGDH activity or deficient in the activity can be selected.

A method for preparing the α KGDH gene deficient strain of *Brevibacterium lactofermentum* by utilizing homologous recombination is described in detail in WO 95/34672. Similar methods can be applied to the other microorganisms.

Further, techniques such as cloning of genes and cleavage and ligation of DNA, transformation and so forth are described in detail in Molecular Cloning, 2nd Edition, Cold Spring Harbor Press, 1989 and so forth.

As a specific example of a mutant strain deficient in

α KGDH activity or with decreased α KGDH activity obtained as described above, there can be mentioned *Enterobacter agglomerans* AJ13356. *Enterobacter agglomerans* AJ13356 was deposited at the National Institute of Bioscience and
5 Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry (postal code: 305-8566, 1-3, Higashi 1-chome, Tsukuba-shi, Ibaraki, Japan) on February 19, 1998 and received an accession number of FERM P-16645. It was then transferred
10 to an international deposition under the provisions of Budapest Treaty on January 11, 1999 and received an accession number of FERM BP-6615. The *Enterobacter agglomerans* AJ13356 is deficient in α KGDH activity as a result of disruption of the α KGDH-E1 subunit gene (*sucA*).

15 When *Enterobacter agglomerans*, an example of the microorganism used in the present invention, is cultured in a medium containing a saccharide, a viscous material is extracellularly secreted, resulting in low operation efficiency. Therefore, when *Enterobacter agglomerans*
20 having such a property of secreting the viscous material is used, it is preferable to use a mutant strain that secretes less the viscous material compared with a wild strain. Examples of mutagenesis methods include, for example,
25 method utilizing treatment with a mutagenic agent such as N-methyl-N'-nitro-N-nitrosoguanidine and so forth. A mutant strain with decreased secretion of the viscous material can be selected by inoculating mutagenized

bacterial cells in a medium containing a saccharide, for example, LB medium plate containing 5 g/L of glucose, culturing them with tilting the plate about 45 degrees and selecting a colony which does not show flowing down of liquid.

5 In the present invention, impartation or enhancement of L-glutamic acid-producing ability and impartation of other favorable properties such as mutation for less viscous material secretion described above can be carried out in an arbitrary order.

10 By culturing the microorganism of the present invention in a liquid medium of which pH is adjusted to a pH at which L-glutamic acid is precipitated, L-glutamic acid can be produced and accumulated with precipitating it in the medium. L-Glutamic acid can also be precipitated by
15 starting the culture at a neutral pH and then ending it at a pH at which L-glutamic acid is precipitated.

The pH at which L-glutamic acid is precipitated means one at which L-glutamic acid is precipitated when the microorganism produces and accumulates L-glutamic acid.

20 As the aforementioned medium, a usual nutrient medium containing a carbon source, a nitrogen source, mineral salts and organic trace nutrients such as amino acids and vitamins as required can be used so long as pH is adjusted to a pH at which L-glutamic acid is precipitated. Either a
25 synthetic medium or a natural medium can be used. The carbon source and the nitrogen source used in the medium can be any ones so long as they can be used by the cultured strain.

As the carbon source, saccharides such as glucose,

glycerol, fructose, sucrose, maltose, mannose, galactose, starch hydrolysate and molasses are used. In addition, organic acids such as acetic acid and citric acid may be used each alone or in combination with another carbon source.

5 As the nitrogen source, ammonia, ammonium salts such as ammonium sulfate, ammonium carbonate, ammonium chloride, ammonium phosphate and ammonium acetate, nitrates and so forth are used.

10 As the organic trace nutrients, amino acids, vitamins, fatty acids, nucleic acids, those containing these substances such as peptone, casamino acid, yeast extract and soybean protein decomposition products are used. When an auxotrophic mutant strain that requires an amino acid and so forth for metabolization or growth is used, the required
15 nutrient must be supplemented.

As mineral salts, phosphates, magnesium salts, calcium salts, iron salts, manganese salts and so forth are used.

As for the culture method, aeration culture is usually performed with controlling the fermentation temperature to
20 be 20 to 42°C and pH to be 3 to 5, preferably 4 to 5, more preferably 4 to 4.7, particularly preferably 4 to 4.5. Thus, after about 10 hours to 4 days of culture, a substantial amount of L-glutamic acid is accumulated in the culture. Accumulated L-glutamic acid exceeding the amount
25 corresponding to the saturation concentration is precipitated in the medium.

After completion of the culture, L-glutamic acid precipitated in the culture can be collected by

centrifugation, filtration or the like. L-Glutamic acid dissolved in the medium can be collected according to known methods. For example, the L-glutamic acid can be isolated by concentrating the culture broth to crystallize it or
5 isolated by ion exchange chromatography or the like. L-Glutamic acid precipitated in the culture broth may be isolated together with L-glutamic acid that have been dissolved in the medium after it is crystallized.

According to the method of the present invention,
10 L-glutamic acid exceeding the amount corresponding to the saturation concentration is precipitated, and the concentration of L-glutamic acid dissolved in the medium is maintained at a constant level. Therefore, influence of L-glutamic acid at a high concentration on microorganisms
15 can be reduced. Accordingly, it becomes possible to breed a microorganism having further improved L-glutamic acid-producing ability. Further, since L-glutamic acid is precipitated as crystals, acidification of the culture broth by accumulation of L-glutamic acid is suppressed, and
20 therefore the amount of alkali used for maintaining pH of the culture can significantly be reduced.

EXAMPLES

Hereafter, the present invention will be more
25 specifically explained with reference to the following examples.

<1> Screening of microorganism having L-glutamic acid resistance in acidic environment

Screening of a microorganism having L-glutamic acid resistance in an acidic environment was performed as follows. Each of about 500 samples obtained from nature including soil, fruits, plant bodies, river water in an amount of 1 g was suspended in 5 mL of sterilized water, of which 200 μ L was coated on 20 mL of solid medium of which pH was adjusted to 4.0 with HCl. The composition of the medium was as follows: 3 g/L of glucose, 1 g/L of $(\text{NH}_4)_2\text{SO}_4$, 0.2 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.5 g/L of KH_2PO_4 , 0.2 g/L of NaCl, 0.1 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.01 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.72 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 0.64 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.72 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.4 mg/L of boric acid, 1.2 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 50 μ g/L of biotin, 50 μ g/L of calcium pantothenate, 50 μ g/L of folic acid, 50 μ g/L of inositol, 50 μ g/L of niacin, 50 μ g/L of p-aminobenzoic acid, 50 μ g/L of pyridoxine hydrochloride, 50 μ g/L of riboflavin, 50 μ g/L of thiamine hydrochloride, 50 mg/L of cycloheximide, 20 g/L of agar.

The media plated on which the above samples were plated were incubated at 28°C, 37°C or 50°C for 2 to 4 days and 378 strains each forming a colony were obtained.

Subsequently, each of the strains obtained as described above was inoculated in a test tube of 16.5 cm in length and 14 mm in diameter containing 3 mL of liquid medium (adjusted to pH 4.0 with HCl) containing a saturation concentration of L-glutamic acid and cultured at 28°C, 37°C or 50°C for 24 hours to 3 days with shaking. Then, the grown strains were selected. The composition of the aforementioned medium was follows: 40 g/L of glucose, 20 g/L

of $(\text{NH}_4)_2\text{SO}_4$, 0.5 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/L of KH_2PO_4 , 0.5 g/L of NaCl, 0.25 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.72 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 0.64 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.72 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.4 mg/L of boric acid, 1.2 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 2 g/L of yeast extract.

Thus, 78 strains of microorganisms having L-glutamic acid resistance in an acidic environment were successfully obtained.

<2> Selection of strains with superior growth rate in acidic environment from microorganisms having L-glutamic acid resistance

The various microorganisms having L-glutamic acid resistance in an acidic environment obtained as described above were each inoculated into a test tube of 16.5 cm in length and 14 mm in diameter containing 3 mL of medium (adjusted to pH 4.0 with HCl) obtained by adding 20 g/L of glutamic acid and 2 g/L of glucose to M9 medium (Sambrook, J., Fritsh, E.F. and Maniatis, T., "Molecular Cloning", Cold Spring Harbor Laboratory Press, 1989), and the turbidity of the medium was measured in the time course to select strains with a favorable growth rate. As a result, as a strain showing favorable growth, the AJ13355 strain was obtained from a soil in Iwata-shi, Shizuoka, Japan. This strain was determined as *Enterobacter agglomerans* based on its bacteriological properties described above.

<3> Acquisition of strain with less viscous material

secretion from *Enterobacter agglomerans* AJ13355 strain

Since the *Enterobacter agglomerans* AJ13355 strain extracellularly secretes a viscous material when cultured in a medium containing a saccharide, operation efficiency is not favorable. Therefore, a strain with less viscous material secretion was obtained by the ultraviolet irradiation method (Miller, J.H. et al., "A Short Course in Bacterial Genetics; Laboratory Manual", p.150, Cold Spring Harbor Laboratory Press, 1992).

10 The *Enterobacter agglomerans* AJ13355 strain was irradiated with ultraviolet ray for 2 minutes at the position 60 cm away from a 60-W ultraviolet lamp and cultured in LB medium overnight to fix mutation. The mutagenized strain was diluted and inoculated in LB medium containing 5 g/L of glucose and 20 g/L of agar so that about 100 colonies per plate would emerge and cultured at 30°C overnight with tilting the plate about 45 degrees, and then 20 colonies showing no flowing down of the viscous material were selected.

20 As a strain satisfying conditions that no revertant emerged even after 5 times of subculture in LB medium containing 5 g/L of glucose and 20 g/L of agar, and that there should be observed growth equivalent to the parent strain in LB medium, LB medium containing 5 g/L of glucose and M9 medium (Sambrook, J. et al., Molecular Cloning, 2nd Edition, Cold Spring Harbor Press, 1989) to which 20 g/L of L-glutamic acid and 2 g/L of glucose were added and of which pH was adjusted to 4.5 with HCl, SC17 strain was selected from the

strains selected above.

<4> Construction of glutamic acid-producing bacterium from *Enterobacter agglomerans* SC17 strain

- 5 (1) Preparation of α KGDH deficient strain from *Enterobacter agglomerans* SC17 strain

A strain deficient in α KGDH and with enhanced L-glutamic acid biosynthetic system was prepared from the *Enterobacter agglomerans* SC17 strain.

- 10 (i) Cloning of α KGDH gene (hereafter, referred to as "*sucAB*") of *Enterobacter agglomerans* AJ13355 strain

The *sucAB* gene of the *Enterobacter agglomerans* AJ13355 strain was cloned by selecting a DNA fragment complementing the acetic acid-unassimilating property of the α KGDH-E1 subunit gene (hereafter, referred to as "*sucA*") deficient strain of *Escherichia coli* from chromosomal DNA of the *Enterobacter agglomerans* AJ13355 strain.

The chromosomal DNA of the *Enterobacter agglomerans* AJ13355 strain was isolated by a method usually employed when chromosomal DNA is extracted from *Escherichia coli* (Text for Bioengineering Experiments, Edited by the Society for Bioscience and Bioengineering, Japan, pp.97-98, Baifukan, 1992). The pTWV228 (resistant to ampicillin) used as a vector was commercially available one from Takara Shuzo Co., Ltd.

The chromosomal DNA of the AJ13355 strain digested with *Eco*T221 and pTWV228 digested with *Pst*I were ligated by using T4 ligase and used to transform the *sucA* deficient

Escherichia coli JRG465 strain (Herbert, J. et al., Mol. Gen. Genetics, 105, 182 (1969)). A strain growing in an acetate minimal medium was selected from the transformant strains obtained above, and a plasmid was extracted from it and designated as pTWVEK101. The *Escherichia coli* JRG465 strain harboring pTWVEK101 recovered auxotrophy for succinic acid or L-lysine and L-methionine besides the acetic acid-assimilating property. This suggests that pTWVEK101 contains the *sucA* gene of *Enterobacter agglomerans*.

Fig. 1 shows the restriction map of a DNA fragment derived from *Enterobacter agglomerans* in pTWVEK101. The determined nucleotide sequence of the hatched portion in Fig. 1 is shown as SEQ ID NO: 1. In this sequence, nucleotide sequences considered to be two full length ORFs and two nucleotide sequences considered to be partial sequences of the ORFs were found. SEQ ID NOS: 2 to 5 show amino acid sequences that can be encoded by these ORFs or partial sequences in an order from the 5' end. As a result of homology search for these, it was revealed that the portion of which nucleotide sequences were determined contained a 3'-end partial sequence of the succinate dehydrogenase iron-sulfur protein gene (*sdhB*), full length *sucA* and α KGDH-E2 subunit gene (*sucB*), and 5'-end partial sequence of the succinyl CoA synthetase β subunit gene (*sucC*). The results of comparison of the amino acid sequences deduced from these nucleotide sequences with those derived from *Escherichia coli* (Eur. J. Biochem., 141, pp.351-359 (1984); Eur. J. Biochem., 141, pp.361-374 (1984); Biochemistry, 24, pp.6245-6252 (1985))

are shown in Figs. 2 to 5. Thus, the amino acid sequences each showed very high homology. In addition, it was found that a cluster of *sdhB-sucA-sucB-sucC* was constituted on the chromosome of *Enterobacter agglomerans* as in *Escherichia coli* (Eur. J. Biochem., 141, pp.351-359 (1984); Eur. J. Biochem., 141, pp.361-374 (1984); Biochemistry, 24, pp.6245-6252 (1985)).

(ii) Acquisition of α KGDH deficient strain derived from *Enterobacter agglomerans* SC17 strain

The homologous recombination was performed by using the *sucAB* gene of *Enterobacter agglomerans* obtained as described above to obtain an α KGDH deficient strain of *Enterobacter agglomerans*.

After pTWVEK101 was digested with *Sph*I to excise a fragment containing *sucA*, the fragment was blunt-ended with Klenow fragment (Takara Shuzo Co., Ltd.) and ligated with pBR322 digested with *Eco*RI and blunt-ended with Klenow fragment, by using T4 DNA ligase (Takara Shuzo Co., Ltd.). The obtained plasmid was digested at the restriction enzyme *Bgl*III recognition site positioned substantially at the center of *sucA* by using this enzyme, blunt-ended with Klenow fragment, and then ligated again by using T4 DNA ligase. It was considered that the *sucA* gene did not function because a frameshift mutation was introduced into *sucA* of the plasmid newly constructed through the above procedure.

The plasmid constructed as described above was digested with a restriction enzyme *Apa*LI, and subjected to

agarose gel electrophoresis to recover a DNA fragment containing *sucA* into which the frameshift mutation was introduced and a tetracycline resistance gene derived from pBR322. The recovered DNA fragment was ligated again by
 5 using T4 DNA ligase to construct a plasmid for disrupting the α KGDH gene.

The plasmid for disrupting the α KGDH gene obtained as described above was used to transform the *Enterobacter agglomerans* SC17 strain by electroporation (Miller, J.H.,
 10 "A Short Course in Bacterial Genetics; Handbook", p.279, Cold Spring Harbor Laboratory Press, U.S.A., 1992), and a strain wherein *sucA* on the chromosome was replaced with a mutant type one by homologous recombination of the plasmid was obtained by using the tetracycline resistance as an indicator.
 15 The obtained strain was designated as SC17*sucA* strain.

In order to confirm that the SC17*sucA* strain was deficient in the α KGDH activity, the enzyme activity was measured by the method of Reed et al. (Reed, L.J. and Mukherjee, B.B., Methods in Enzymology, 13, pp.55-61,
 20 (1969)) by using cells of the strain cultured in LB medium until the logarithmic growth phase. As a result, α KGDH activity of 0.073 (Δ ABS/min/mg protein) was detected from the SC17 strain, whereas no α KGDH activity was detected from the SC17*sucA* strain, and thus it was confirmed that the *sucA*
 25 was deficient as purposed.

(2) Enhancement of L-glutamic acid biosynthetic system of *Enterobacter agglomerans* SC17*sucA* strain

Subsequently, a citrate synthase gene, a phosphoenolpyruvate carboxylase gene and a glutamate dehydrogenase gene derived from *Escherichia coli* were introduced into the SC17sucA strain.

5

(i) Preparation of plasmid having *gltA* gene, *ppc* gene and *gdhA* gene derived from *Escherichia coli*

The procedures of preparing a plasmid having a *gltA* gene, a *ppc* gene and a *gdhA* gene will be explained by referring to Figs. 6 and 7.

A plasmid having a *gdhA* gene derived from *Escherichia coli*, pBRGDH (Japanese Patent Application Laid-open No. 7-203980), was digested with *Hind*III and *Sph*I, the both ends were blunt-ended by the T4 DNA polymerase treatment, and then the DNA fragment having the *gdhA* gene was purified and recovered. Separately, a plasmid having a *gltA* gene and a *ppc* gene derived from *Escherichia coli*, pMWCP (WO 97/08294), was digested with *Xba*I, and then the both ends were blunt-ended by using T4 DNA polymerase. This was mixed with the above purified DNA fragment having the *gdhA* gene and ligated by using T4 ligase to obtain a plasmid pMWCPG, which corresponded to pMWCP further containing the *gdhA* gene (Fig. 6).

At the same time, the plasmid pVIC40 (Japanese Patent Application Laid-open No. 8-047397) having the replication origin of the broad host spectrum plasmid RSF1010 was digested with *Not*I, treated with T4 DNA polymerase and digested with *Pst*I. pBR322 was digested with *Eco*T14I,

treated with T4 DNA polymerase and digested with *Pst*I. The both products were mixed and ligated by using T4 ligase to obtain a plasmid RSF-Tet having the replication origin of RSF1010 and a tetracycline resistance gene (Fig. 7).

5 Subsequently, pMWCPG was digested with *Eco*RI and *Pst*I, and a DNA fragment having the *gltA* gene, the *ppc* gene and the *gdhA* gene was purified and recovered. RSF-Tet was similarly digested with *Eco*RI and *Pst*I, and a DNA fragment having the replication origin of RSF1010 was purified and
10 recovered. The both products were mixed and ligated by using T4 ligase to obtain a plasmid RSFCPG, which corresponded to RSF-Tet containing the *gltA* gene, the *ppc* gene and the *gdhA* gene (Fig. 8). It was confirmed that the obtained plasmid RSFCPG expressed the *gltA* gene, the *ppc* gene and the *gdhA*
15 gene, by the complementation of the auxotrophy of the *gltA*, *ppc* or *gdhA* gene deficient strain derived from *Escherichia coli* and measurement of each enzyme activity.

(ii) Preparation of plasmid having *gltA* gene derived from
20 *Brevibacterium lactofermentum*

 A plasmid having the *gltA* gene derived from *Brevibacterium lactofermentum* was constructed as follows. PCR was performed by using the primer DNAs having the nucleotide sequences represented by SEQ ID NOS: 6 and 7, which
25 were prepared based on the nucleotide sequence of the *Corynebacterium glutamicum gltA* gene (Microbiology, 140, pp.1817-1828 (1994)), and chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 as a template to obtain a *gltA* gene

fragment of about 3 kb. This fragment inserted into a plasmid pHSG399 (purchased from Takara Shuzo Co., Ltd.) digested with *Sma*I to obtain a plasmid pHSGCB (Fig. 9). Subsequently, pHSGCB was digested with *Hind*III, and the excised *gltA* gene fragment of about 3 kb was inserted into a plasmid pSTV29 (purchased from Takara Shuzo Co., Ltd.) digested with *Hind*III to obtain a plasmid pSTVCB (Fig. 9). It was confirmed that the obtained plasmid pSTVCB expressed the *gltA* gene, by measuring the enzyme activity in the *Enterobacter agglomerans* AJ13355 strain.

(iii) Introduction of RSFCPG and pSTVCB into SC17sucA strain

The *Enterobacter agglomerans* SC17sucA strain was transformed with RSFCPG by electroporation to obtain a transformant SC17sucA/RSFCPG strain having tetracycline resistance. Further, the SC17sucA/RSFCPG strain was transformed with pSTVCB by electroporation to obtain a transformant SC17sucA/RSFCPG+pSTVCB strain having chloramphenicol resistance.

<4> Acquisition of strain with improved resistance to L-glutamic acid in low pH environment

A strain with improved resistance to L-glutamic acid at a high concentration in a low pH environment (hereafter, also referred to as "high-concentration Glu-resistant strain at low pH") was isolated from the *Enterobacter agglomerans* SC17sucA/RSFCPG+pSTVCB strain.

The SC17sucA/RSFCPG+pSTVCB strain was cultured

overnight at 30°C in LBG medium (10 g/L of tryptone, 5 g/L of yeast extract, 10 g/L of NaCl, 5 g/L of glucose), and the cells washed with saline was appropriately diluted and plated on an M9-E medium (4 g/L of glucose, 17 g/L of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, 3 g/L of KH_2PO_4 , 0.5 g/L of NaCl, 1 g/L of NH_4Cl , 10 mM of MgSO_4 , 10 μM of CaCl_2 , 50 mg/L of L-lysine, 50 mg/L of L-methionine, 50 mg/L of DL-diaminopimelic acid, 25 mg/L of tetracycline, 25 mg/L of chloramphenicol, 30 g/L of L-glutamic acid, adjusted to pH 4.5 with aqueous ammonia) plate.

10 The colony emerged after culture at 32°C for 2 days was obtained as a high-concentration Glu-resistant strain at low pH.

For the obtained strain, growth level in M9-E liquid medium was measured and L-glutamic acid-producing ability was tested in a 50-ml volume large test tube containing 5 ml of L-glutamic acid production test medium (40 g/L of glucose, 20 g/L of $(\text{NH}_4)_2\text{SO}_4$, 0.5 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/L of KH_2PO_4 , 0.5 g/L of NaCl, 0.25 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.72 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 0.64 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.72 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.4 mg/L of boric acid, 1.2 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 2 g/L of yeast extract, 200 mg/L of L-lysine hydrochloride, 200 mg/L of L-methionine, 200 mg/L of DL- α,ϵ -diaminopimelic acid, 25 mg/L of tetracycline hydrochloride, 25 mg/L of chloramphenicol).

25 A strain that exhibited the best growth level and the same L-glutamic acid producing ability as that of its parent strain, the SC17/RSFCPG+pSTVCB strain, was designated as *Enterobacter agglomerans* AJ13601. The AJ13601 strain was

deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry (postal code: 305-8566, 1-3, Higashi 1-chome, Tsukuba-shi, Ibaraki, Japan) on August 18, 1999 and received an accession number of FERM P-17516. It was then transferred to an international deposition under the provisions of Budapest Treaty on July 6, 2000 and received an accession number of FERM BP-7207.

10

<5> Culture of *Enterobacter agglomerans* AJ13601 strain for L-glutamic acid production (1)

The *Enterobacter agglomerans* AJ13601 strain was inoculated into a 1-L jar fermenter containing 300 ml of medium containing 40 g/L of glucose, 20 g/L of $(\text{NH}_4)_2\text{SO}_4$, 0.5 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/L of KH_2PO_4 , 0.5 g/L of NaCl, 0.25 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.72 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 0.64 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.72 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.4 mg/L of boric acid, 1.2 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 2 g/L of yeast extract, 200 mg/L of L-lysine hydrochloride, 200 mg/L of L-methionine, 200 mg/L of DL- α,ϵ -diaminopimelic acid, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol, and cultured at 34°C and pH 6.0 for 14 hours. The culture pH was controlled by introducing ammonia gas into the medium.

The culture obtained as described above was centrifuged at 5000 rpm for 10 minutes, and the collected cells were inoculated into a 1-L jar fermenter containing

300 ml of medium containing 40 g/L of glucose, 5 g/L of $(\text{NH}_4)_2\text{SO}_4$, 1.5 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 6 g/L of KH_2PO_4 , 1.5 g/L of NaCl , 0.75 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.06 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.06 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 2.16 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 1.92 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 2.16 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 1.2 mg/L of boric acid, 3.6 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 6 g/L of yeast extract, 600 mg/L of L-lysine hydrochloride, 600 mg/L of L-methionine, 600 mg/L of DL- α, ϵ -diaminopimelic acid, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol and cultured at 34°C and pH 4.5 to perform culture for L-glutamic acid production. The culture pH was controlled by introducing ammonia gas into the medium. As the initially added glucose was depleted, 600 g/L of glucose was continuously added.

As a result of the culture for L-glutamic acid production performed for 50 hours as described above, a substantial amount of L-glutamic acid crystals were precipitated in the jar fermenter. Table 1 shows the concentration of L-glutamic acid dissolved in the culture broth at that time and the L-glutamic acid concentration measured by dissolving the crystals in 2 M potassium hydroxide. L-Glutamic acid crystals were collected from the culture by decantation after the culture was left stood.

Table 1

Concentration of L-glutamic acid dissolved in culture broth	51 g/L
Amount of L-glutamic acid precipitated as crystals	67 g/L
Concentration of L-glutamic acid measured by dissolving crystals	118 g/L

<6> Culture of *Enterobacter agglomerans* AJ13601 strain for L-glutamic acid production (2)

The following experiment was performed in order to confirm that the *Enterobacter agglomerans* AJ13601 strain still had L-glutamic acid-producing ability even under the condition that L-glutamic acid crystals were present.

The *Enterobacter agglomerans* AJ13601 strain was inoculated into a 1-L jar fermenter containing 300 ml of medium containing 40 g/L of glucose, 20 g/L of $(\text{NH}_4)_2\text{SO}_4$, 0.5 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/L of KH_2PO_4 , 0.5 g/L of NaCl, 0.25 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.02 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.72 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 0.64 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.72 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.4 mg/L of boric acid, 1.2 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 2 g/L of yeast extract, 200 mg/L of L-lysine hydrochloride, 200 mg/L of L-methionine, 200 mg/L of DL- α,ϵ -diaminopimelic acid, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol, and cultured at 34°C at pH 6.0 for 14 hours. The culture pH was controlled by bubbling the medium with ammonia gas. The culture obtained as described above was centrifuged at 5000 rpm for 10 minutes, and then the collected cells were cultured in a medium where L-glutamic acid was present as crystals. The used medium contained 40 g/L of glucose, 5 g/L of $(\text{NH}_4)_2\text{SO}_4$, 1.5 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 6 g/L of KH_2PO_4 , 1.5 g/L of NaCl, 0.75 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.06 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.06 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 2.16 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 1.92 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 2.16 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 1.2 mg/L of boric acid, 3.6 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 6 g/L of yeast extract, 600

mg/L of L-lysine hydrochloride, 600 mg/L of L-methionine, 600 mg/L of DL- α,ϵ -diaminopimelic acid, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol and L-glutamic acid crystals were added to 40 g/L. The cells
5 were inoculated in a 1-L jar fermenter containing 300 ml of this medium and cultured at 34°C and pH 4.3 to perform culture for L-glutamic acid production. The culture pH was controlled by introducing ammonia gas into the medium. As the initially added glucose was depleted, 600 g/L of glucose
10 was continuously added. In this medium, only 39 g/L of the added L-glutamic acid was dissolved at pH 4.3 and the remaining 1 g/L was present as crystals.

As a result of the culture for L-glutamic acid production performed for 53 hours as described above, a
15 substantial amount of L-glutamic acid crystals were precipitated in the jar fermenter. Table 2 shows the concentration of L-glutamic acid dissolved in the culture broth, the amount of L-glutamic acid present as crystals at that time and the L-glutamic acid concentration measured by
20 dissolving the crystals in 2 M KOH solution. L-Glutamic acid crystals were collected from the culture by decantation after the culture was left stood. The results showed that the *Enterobacter agglomerans* AJ13601 strain accumulated L-glutamic acid and precipitated crystals thereof even under
25 the condition that L-glutamic acid crystals were present.

Table 2

Concentration of L-glutamic acid dissolved in culture broth	39 g/L
Amount of L-glutamic acid precipitated as crystals	119 g/L
Concentration of L-glutamic acid measured by dissolving crystals	158 g/L
Amount of L-glutamic acid crystals newly produced by main culture	118 g/L

<7> Culture of *Enterobacter agglomerans* AJ13601 strain for L-glutamic acid production (3)

5 The *Enterobacter agglomerans* AJ13601 strain can grow not only at an acidic pH, but also at a neutral pH. Therefore, it was confirmed as follows that L-glutamic acid crystals could also be precipitated by starting the culture at a neutral pH and allowing production of L-glutamic acid during
10 the culture so that pH of the culture should spontaneously be lowered.

Cells of one plate (8.5 cm in diameter) of the *Enterobacter agglomerans* AJ13601 strain, cultured on LBG agar medium (10 g/ of L tryptone, 5 g/L of yeast extract,
15 10 g/L of NaCl, 5 g/L of glucose, 15 g/L of agar) containing 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol at 30°C for 14 hours, were inoculated into a 1-L jar fermenter containing 300 ml of medium containing
20 40 g/L of glucose, 5 g/L of $(\text{NH}_4)_2\text{SO}_4$, 1.5 g/L of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 6 g/L of KH_2PO_4 , 1.5 g/L of NaCl, 0.75 g/L of $\text{CaCl}_2 \cdot 7\text{H}_2\text{O}$, 0.06 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.06 g/L of $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 2.16 mg/L of $\text{ZnSO}_4 \cdot 2\text{H}_2\text{O}$, 1.92 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 2.16 mg/L of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 1.2 mg/L of boric acid, 3.6 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 6 g/L of yeast extract,

600 mg/L of L-lysine hydrochloride, 600 mg/L of L-methionine, 600 mg/L of DL- α,ϵ -diaminopimelic acid, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol and the culture was started at 34°C and pH 7.0. The culture
5 pH was controlled by introducing ammonia gas into the medium. As the initially added glucose was depleted, 600 g/L of glucose was continuously added.

As L-glutamic acid is accumulated, pH lowers spontaneously. The amount of the introduced ammonia gas was
10 adjusted so that pH should be gradually lowered from 7.0 to 4.5 during the period between 15 hours and 24 hours after the start of the culture, and 24 hours after the start of the culture, pH became 4.5. Afterward, cultivation was continued for 12 hours.

15 As a result of the culture for L-glutamic acid production conducted for 36 hours as described above, a substantial amount of L-glutamic acid crystals were precipitated in the jar fermenter. Table 3 shows the concentration of L-glutamic acid dissolved in the culture
20 broth, the amount of L-glutamic acid present as crystals at that time and the L-glutamic acid concentration measured by dissolving the crystals in 2 M KOH solution. L-Glutamic acid crystals were collected from the culture by decantation after the culture was left stood.

Table 3

Concentration of L-glutamic acid dissolved in culture broth	45 g/L
Amount of L-glutamic acid precipitated as crystals	31 g/L
Concentration of L-glutamic acid measured by dissolving crystals	76 g/L

SEQUENCE LISTING

<110> Hiroshi Izui
Mika Moriya
Seiko Hirano
Yoshihiko Hara
Hisao Ito
Kazuhiko Matsui

<120> Method for producing L-glutamic acid by fermentation accompanied by precipitation

<130>

<150> JP 11-234806

<151> 1999-08-20

<150> JP 2000-78771

<151> 2000-03-21

<160> 7

<170> PatentIn Ver. 2.0

<210> 1

<211> 4556

<212> DNA

<213> Enterobacter agglomerans

<220>

<221> CDS

<222> (2)..(121)

<220>

<221> CDS

<222> (322)..(3129)

<220>

<221> CDS

<222> (3145)..(4368)

<220>

<221> CDS

<222> (4437)..(4556)

<400> 1

```

t gca ttc agc gtt ttc cgc tgt cac agc atc atg aac tgt gta agt gtt 49
Ala Phe Ser Val Phe Arg Cys His Ser Ile Met Asn Cys Val Ser Val
      1           5           10           15
tgt cct aaa ggg cta aac ccg acg cgc gct atc ggc cac att aag tcg 97
Cys Pro Lys Gly Leu Asn Pro Thr Arg Ala Ile Gly His Ile Lys Ser
                20                25                30
atg ctg ctg caa cgc agc gcg tagttatacc accggaacc tcaggttccc 148
Met Leu Leu Gln Arg Ser Ala

```

35
 ggtattttac ggaagcctct gtaaacgcgg tccaaccac gtttacaag gttcccttac 208
 gggccgggcg cgcgctgcgc acagtgcctg tatcgctgaa ctactacgg caaaccgcga 268
 aagcggcaac aaatgaaacc tcaaaaaagc ataacattgc ttaagggatc aca atg 324
 Met
 1
 cag aac agc gcg atg aag ccc tgg ctg gac tcc tcc tgg ctg gcc ggc 372
 Gln Asn Ser Ala Met Lys Pro Trp Leu Asp Ser Ser Trp Leu Ala Gly
 5 10 15
 gcg aat cag tct tac ata gag caa ctc tat gag gat ttc ctg acc gat 420
 Ala Asn Gln Ser Tyr Ile Glu Gln Leu Tyr Glu Asp Phe Leu Thr Asp
 20 25 30
 cct gac tct gtg gat gca gtg tgg cgc tcg atg ttc caa cag tta cca 468
 Pro Asp Ser Val Asp Ala Val Trp Arg Ser Met Phe Gln Gln Leu Pro
 35 40 45
 ggc acg gga gtg aaa cct gag cag ttc cac tcc gca act cgc gaa tat 516
 Gly Thr Gly Val Lys Pro Glu Gln Phe His Ser Ala Thr Arg Glu Tyr
 50 55 60 65
 ttc cgt cgc ctg gcg aaa gac gca tct cgt tac acc tcc tca gtt acc 564
 Phe Arg Arg Leu Ala Lys Asp Ala Ser Arg Tyr Thr Ser Ser Val Thr
 70 75 80
 gat ccg gca acc aac tcc aaa caa gtg aaa gtg ctg cag ctg att aac 612
 Asp Pro Ala Thr Asn Ser Lys Gln Val Lys Val Leu Gln Leu Ile Asn
 85 90 95
 gcg ttt cgt ttc cgc gga cat cag gaa gca aat ctc gat ccg ctt ggc 660
 Ala Phe Arg Phe Arg Gly His Gln Glu Ala Asn Leu Asp Pro Leu Gly
 100 105 110
 ctg tgg aaa cag gac cgc gtt gcc gat ctc gat cct gcc ttt cac gat 708
 Leu Trp Lys Gln Asp Arg Val Ala Asp Leu Asp Pro Ala Phe His Asp
 115 120 125
 ctg acc gac gcc gat ttt cag gaa agc ttt aac gta ggt tct ttt gcc 756
 Leu Thr Asp Ala Asp Phe Gln Glu Ser Phe Asn Val Gly Ser Phe Ala
 130 135 140 145
 att ggc aaa gaa acc atg aag ctg gcc gat ctg ttc gac gcg ctg aag 804
 Ile Gly Lys Glu Thr Met Lys Leu Ala Asp Leu Phe Asp Ala Leu Lys
 150 155 160
 cag acc tac tgt ggc tcg att ggt gca gag tat atg cac atc aat aac 852
 Gln Thr Tyr Cys Gly Ser Ile Gly Ala Glu Tyr Met His Ile Asn Asn
 165 170 175
 acc gaa gag aaa cgc tgg atc cag cag cgt atc gaa tcc ggt gcg agc 900
 Thr Glu Glu Lys Arg Trp Ile Gln Gln Arg Ile Glu Ser Gly Ala Ser
 180 185 190
 cag acg tca ttc agt ggc gaa gag aaa aaa ggt ttc ctg aaa gag ctg 948
 Gln Thr Ser Phe Ser Gly Glu Glu Lys Lys Gly Phe Leu Lys Glu Leu
 195 200 205
 acc gcg gca gaa ggg ctg gaa aaa tat ctg ggc gcg aaa ttc ccg ggt 996
 Thr Ala Ala Glu Gly Leu Glu Lys Tyr Leu Gly Ala Lys Phe Pro Gly
 210 215 220 225
 gca aaa cgt ttc tcg ctg gaa ggc ggt gat gcg ctg gtg ccg atg ctg 1044
 Ala Lys Arg Phe Ser Leu Glu Gly Gly Asp Ala Leu Val Pro Met Leu
 230 235 240
 cgc gag atg att cgt cat gcg ggc aaa agc ggc aca cgt gaa gtg gta 1092
 Arg Glu Met Ile Arg His Ala Gly Lys Ser Gly Thr Arg Glu Val Val
 245 250 255

ctg	ggg	atg	gcg	cac	cgt	ggc	cgt	ctt	aac	gta	ctg	att	aac	gta	ctg	1140
Leu	Gly	Met	Ala	His	Arg	Gly	Arg	Leu	Asn	Val	Leu	Ile	Asn	Val	Leu	
	260					265					270					
ggt	aaa	aag	cca	cag	gat	ctg	ttc	gac	gaa	ttc	tcc	ggt	aaa	cac	aaa	1188
Gly	Lys	Lys	Pro	Gln	Asp	Leu	Phe	Asp	Glu	Phe	Ser	Gly	Lys	His	Lys	
	275					280					285					
gag	cat	ctg	ggc	acc	ggt	gat	gtg	aag	tat	cac	atg	ggc	ttc	tct	tcg	1236
Glu	His	Leu	Gly	Thr	Gly	Asp	Val	Lys	Tyr	His	Met	Gly	Phe	Ser	Ser	
	290				295					300					305	
gat	att	gaa	acc	gaa	ggt	ggt	ctg	gtg	cat	ctg	gcg	ctg	gcg	ttt	aac	1284
Asp	Ile	Glu	Thr	Glu	Gly	Gly	Leu	Val	His	Leu	Ala	Leu	Ala	Phe	Asn	
				310					315					320		
ccg	tct	cac	ctg	gaa	att	gtc	agc	ccg	gtg	gtc	atg	gga	tcg	gta	cgt	1332
Pro	Ser	His	Leu	Glu	Ile	Val	Ser	Pro	Val	Val	Met	Gly	Ser	Val	Arg	
			325					330					335			
gca	cgt	ctc	gat	cgt	ctg	gcc	gaa	ccg	gtc	agc	aat	aaa	gtg	ttg	cct	1380
Ala	Arg	Leu	Asp	Arg	Leu	Ala	Glu	Pro	Val	Ser	Asn	Lys	Val	Leu	Pro	
			340			345						350				
atc	acc	att	cac	ggt	gat	gcg	gcg	gtg	att	ggt	cag	ggc	gtg	gtt	cag	1428
Ile	Thr	Ile	His	Gly	Asp	Ala	Ala	Val	Ile	Gly	Gln	Gly	Val	Val	Gln	
	355				360						365					
gaa	acc	ctg	aac	atg	tct	cag	gcg	cgc	ggc	tac	gaa	gtg	ggc	ggc	acg	1476
Glu	Thr	Leu	Asn	Met	Ser	Gln	Ala	Arg	Gly	Tyr	Glu	Val	Gly	Gly	Thr	
	370				375					380					385	
gta	cgt	atc	gtc	att	aac	aac	cag	ggt	ggt	ttt	acc	acc	tcc	aac	ccg	1524
Val	Arg	Ile	Val	Ile	Asn	Asn	Gln	Val	Gly	Phe	Thr	Thr	Ser	Asn	Pro	
				390					395					400		
aaa	gat	gcg	cgt	tca	acc	ccg	tac	tgt	act	gac	atc	ggc	aag	atg	gtg	1572
Lys	Asp	Ala	Arg	Ser	Thr	Pro	Tyr	Cys	Thr	Asp	Ile	Gly	Lys	Met	Val	
			405					410					415			
ctg	gca	ccg	att	ttc	cac	gtc	aat	gct	gac	gat	ccg	gaa	gcg	gtg	gcc	1620
Leu	Ala	Pro	Ile	Phe	His	Val	Asn	Ala	Asp	Asp	Pro	Glu	Ala	Val	Ala	
			420					425				430				
ttt	ggt	acc	cgc	ctg	gcg	ctg	gac	tat	cgc	aac	acc	ttc	aaa	cgc	gat	1668
Phe	Val	Thr	Arg	Leu	Ala	Leu	Asp	Tyr	Arg	Asn	Thr	Phe	Lys	Arg	Asp	
	435					440					445					
gtg	ttt	atc	gat	ctg	gtg	tgc	tat	cgc	cgt	cat	ggt	cac	aac	gag	gcg	1716
Val	Phe	Ile	Asp	Leu	Val	Cys	Tyr	Arg	Arg	His	Gly	His	Asn	Glu	Ala	
	450				455					460					465	
gat	gag	cca	agt	gct	acc	cag	ccg	ttg	atg	tac	cag	aaa	atc	aaa	aag	1764
Asp	Glu	Pro	Ser	Ala	Thr	Gln	Pro	Leu	Met	Tyr	Gln	Lys	Ile	Lys	Lys	
				470					475					480		
cat	ccg	acg	ccg	cgt	aaa	att	tac	gcc	gat	cgt	ctg	gaa	ggc	gaa	ggt	1812
His	Pro	Thr	Pro	Arg	Lys	Ile	Tyr	Ala	Asp	Arg	Leu	Glu	Gly	Glu	Gly	
			485					490					495			
gtc	gcg	tcg	cag	gaa	gat	gcc	acc	gag	atg	gtg	aac	ctg	tac	cgc	gat	1860
Val	Ala	Ser	Gln	Glu	Asp	Ala	Thr	Glu	Met	Val	Asn	Leu	Tyr	Arg	Asp	
			500					505					510			
gcg	ctc	gat	gcg	ggc	gaa	tgc	gtg	gtg	ccg	gaa	tgg	cgt	ccg	atg	agc	1908
Ala	Leu	Asp	Ala	Gly	Glu	Cys	Val	Val	Pro	Glu	Trp	Arg	Pro	Met	Ser	
			515			520					525					
ctg	cac	tcc	ttc	acg	tgg	tcg	cct	tat	ctg	aac	cac	gaa	tgg	gat	gag	1956
Leu	His	Ser	Phe	Thr	Trp	Ser	Pro	Tyr	Leu	Asn	His	Glu	Trp	Asp	Glu	
	530				535					540					545	

cct tat ccg gca cag gtt gac atg aaa cgc ctg aag gaa ctg gca ttg 2004
 Pro Tyr Pro Ala Gln Val Asp Met Lys Arg Leu Lys Glu Leu Ala Leu
 550 555 560
 cgt atc agc cag gtc cct gag cag att gaa gtg cag tcg cgc gtg gcc 2052
 Arg Ile Ser Gln Val Pro Glu Gln Ile Glu Val Gln Ser Arg Val Ala
 565 570 575
 aag atc tat aac gat cgc aag ctg atg gcc gaa ggc gag aaa gcg ttc 2100
 Lys Ile Tyr Asn Asp Arg Lys Met Ala Glu Gly Glu Lys Ala Phe
 580 585 590
 gac tgg ggc ggt gcc gag aat ctg gcg tac gcc acg ctg gtg gat gaa 2148
 Asp Trp Gly Gly Ala Glu Asn Leu Ala Tyr Ala Thr Leu Val Asp Glu
 595 600 605
 ggt att ccg gtt cgc ctc tcg ggt gaa gac tcc ggt cgt gga acc ttc 2196
 Gly Ile Pro Val Arg Leu Ser Gly Glu Asp Ser Gly Arg Gly Thr Phe
 610 615 620 625
 ttc cat cgc cac gcg gtc gtg cac aac cag gct aac ggt tca acc tat 2244
 Phe His Arg His Ala Val Val His Asn Gln Ala Asn Gly Ser Thr Tyr
 630 635 640
 acg ccg ctg cac cat att cat aac agc cag ggc gag ttc aaa gtc tgg 2292
 Thr Pro Leu His His Ile His Asn Ser Gln Gly Glu Phe Lys Val Trp
 645 650 655
 gat tcg gtg ctg tct gaa gaa gcg gtg ctg gcg ttt gaa tac ggt tac 2340
 Asp Ser Val Leu Ser Glu Glu Ala Val Leu Ala Phe Glu Tyr Gly Tyr
 660 665 670
 gcc acg gct gag ccg cgc gtg ctg acc atc tgg gaa gcg cag ttt ggt 2388
 Ala Thr Ala Glu Pro Arg Val Leu Thr Ile Trp Glu Ala Gln Phe Gly
 675 680 685
 gac ttt gcc aac ggt gct cag gtg gtg att gac cag ttc atc agc tct 2436
 Asp Phe Ala Asn Gly Ala Gln Val Val Ile Asp Gln Phe Ile Ser Ser
 690 695 700 705
 ggc gaa cag aag tgg ggc cgt atg tgt ggc ctg gtg atg ttg ctg ccg 2484
 Gly Glu Gln Lys Trp Gly Arg Met Cys Gly Leu Val Met Leu Leu Pro
 710 715 720
 cat ggc tac gaa ggt cag gga ccg gaa cac tcc tct gcc cgt ctg gaa 2532
 His Gly Tyr Glu Gly Gln Gly Pro Glu His Ser Ser Ala Arg Leu Glu
 725 730 735
 cgc tat ctg caa ctt tgc gcc gag cag aac atg cag gtt tgc gtc ccg 2580
 Arg Tyr Leu Gln Leu Cys Ala Glu Gln Asn Met Gln Val Cys Val Pro
 740 745 750
 tcg acg ccg gct cag gtg tat cac atg ctg cgc cgt cag gcg ctg cgc 2628
 Ser Thr Pro Ala Gln Val Tyr His Met Leu Arg Arg Gln Ala Leu Arg
 755 760 765
 ggg atg cgc cgt ccg ctg gtg gtg atg tcg ccg aag tcg ctg tta cgc 2676
 Gly Met Arg Arg Pro Leu Val Val Met Ser Pro Lys Ser Leu Leu Arg
 770 775 780 785
 cat cca ctg gcg atc tcg tcg ctg gat gaa ctg gca aac ggc agt ttc 2724
 His Pro Leu Ala Ile Ser Ser Leu Asp Glu Leu Ala Asn Gly Ser Phe
 790 795 800
 cag ccg gcc att ggt gag atc gac gat ctg gat ccg cag ggc gtg aaa 2772
 Gln Pro Ala Ile Gly Glu Ile Asp Asp Leu Asp Pro Gln Gly Val Lys
 805 810 815
 cgc gtc gtg ctg tgc tcc ggt aag gtt tac tac gat ctg ctg gaa cag 2820
 Arg Val Val Leu Cys Ser Gly Lys Val Tyr Tyr Asp Leu Leu Glu Gln
 820 825 830

cgt	cgt	aaa	gac	gag	aaa	acc	gat	gtt	gcc	atc	gtg	cgc	atc	gaa	cag	2868
Arg	Arg	Lys	Asp	Glu	Lys	Thr	Asp	Val	Ala	Ile	Val	Arg	Ile	Glu	Gln	
835						840				845						
ctt	tac	ccg	ttc	ccg	cat	cag	gcg	gta	cag	gaa	gca	ttg	aaa	gcc	tat	2916
Leu	Tyr	Pro	Phe	Pro	His	Gln	Ala	Val	Gln	Glu	Ala	Leu	Lys	Ala	Tyr	
850					855				860						865	
tct	cac	gta	cag	gac	ttt	gtc	tgg	tgc	cag	gaa	gag	cct	ctg	aac	cag	2964
Ser	His	Val	Gln	Asp	Phe	Val	Trp	Cys	Gln	Glu	Glu	Pro	Leu	Asn	Gln	
			870						875					880		
ggc	gcc	tgg	tac	tgt	agc	cag	cat	cat	ttc	cgt	gat	gtc	gtg	ccg	ttt	3012
Gly	Ala	Trp	Tyr	Cys	Ser	Gln	His	His	Phe	Arg	Asp	Val	Val	Pro	Phe	
			885					890					895			
ggc	gcc	acc	ctg	cgt	tat	gca	ggc	cgc	ccg	gca	tcg	gct	tct	ccg	gcc	3060
Gly	Ala	Thr	Leu	Arg	Tyr	Ala	Gly	Arg	Pro	Ala	Ser	Ala	Ser	Pro	Ala	
	900					905						910				
gtg	ggc	tat	atg	tcc	gta	cac	caa	caa	cag	cag	caa	gac	ctg	gtt	aat	3108
Val	Gly	Tyr	Met	Ser	Val	His	Gln	Gln	Gln	Gln	Gln	Asp	Leu	Val	Asn	
	915				920						925					
gac	gca	ctg	aac	gtc	aat	taattaaag	gaaagata	atg	agt	agc	gta	gat				3159
Asp	Ala	Leu	Asn	Val	Asn			Met	Ser	Ser	Val	Asp				
930					935			1				5				
att	ctc	gtt	ccc	gac	ctg	cct	gaa	tcg	gtt	gca	gat	gcc	aca	gta	gca	3207
Ile	Leu	Val	Pro	Asp	Leu	Pro	Glu	Ser	Val	Ala	Asp	Ala	Thr	Val	Ala	
			10					15						20		
acc	tgg	cac	aag	aaa	cca	ggc	gat	gca	gtc	agc	cgc	gat	gaa	gtc	atc	3255
Thr	Trp	His	Lys	Lys	Pro	Gly	Asp	Ala	Val	Ser	Arg	Asp	Glu	Val	Ile	
			25				30						35			
gtc	gaa	att	gaa	act	gac	aaa	gtc	gtg	ctg	gaa	gtg	ccg	gca	tct	gcc	3303
Val	Glu	Ile	Glu	Thr	Asp	Lys	Val	Val	Leu	Glu	Val	Pro	Ala	Ser	Ala	
	40					45						50				
gat	ggc	gtg	ctg	gaa	gcc	gtg	ctg	gaa	gac	gaa	ggg	gca	acc	gtt	acg	3351
Asp	Gly	Val	Leu	Glu	Ala	Val	Leu	Glu	Asp	Glu	Gly	Ala	Thr	Val	Thr	
	55				60					65						
tcc	cgc	cag	atc	ctg	ggc	cgc	ctg	aaa	gaa	ggc	aac	agt	gcg	ggc	aaa	3399
Ser	Arg	Gln	Ile	Leu	Gly	Arg	Leu	Lys	Glu	Gly	Asn	Ser	Ala	Gly	Lys	
	70				75				80						85	
gaa	agc	agt	gcc	aaa	gcg	gaa	agc	aat	gac	acc	acg	cca	gcc	cag	cgt	3447
Glu	Ser	Ser	Ala	Lys	Ala	Glu	Ser	Asn	Asp	Thr	Thr	Pro	Ala	Gln	Arg	
			90					95						100		
cag	aca	gcg	tcg	ctt	gaa	gaa	gag	agc	gat	gcg	ctc	agc	ccg	gcg		3495
Gln	Thr	Ala	Ser	Leu	Glu	Glu	Glu	Ser	Ser	Asp	Ala	Leu	Ser	Pro	Ala	
			105				110					115				
atc	cgt	cgc	ctg	att	gcg	gag	cat	aat	ctt	gac	gct	gcg	cag	atc	aaa	3543
Ile	Arg	Arg	Leu	Ile	Ala	Glu	His	Asn	Leu	Asp	Ala	Ala	Gln	Ile	Lys	
	120					125						130				
ggc	acc	ggc	gta	ggc	gga	cgt	tta	acg	cgt	gaa	gac	gtt	gaa	aaa	cat	3591
Gly	Thr	Gly	Val	Gly	Gly	Arg	Leu	Thr	Arg	Glu	Asp	Val	Glu	Lys	His	
	135					140				145						
ctg	gcg	aac	aaa	ccg	cag	gct	gag	aaa	gcc	gcc	gcg	cca	gcg	gcg	ggc	3639
Leu	Ala	Asn	Lys	Pro	Gln	Ala	Glu	Lys	Ala	Ala	Ala	Pro	Ala	Ala	Gly	
	150				155				160						165	
gca	gca	acg	gct	cag	cag	cct	gtt	gcc	aac	cgc	agc	gaa	aaa	cgt	gtt	3687
Ala	Ala	Thr	Ala	Gln	Gln	Pro	Val	Ala	Asn	Arg	Ser	Glu	Lys	Arg	Val	
				170					175						180	

ccg atg acg cgt tta cgt aag cgc gtc gcg gag cgt ctg ctg gaa gcc 3735
 Pro Met Thr Arg Leu Arg Lys Arg Val Ala Glu Arg Leu Leu Glu Ala
 185 190 195
 aag aac agc acc gcc atg ttg acg acc ttc aac gaa atc aac atg aag 3783
 Lys Asn Ser Thr Ala Met Leu Thr Thr Phe Asn Glu Ile Asn Met Lys
 200 205 210
 ccg att atg gat ctg cgt aag cag tac ggc gat gcg ttc gag aag cgt 3831
 Pro Ile Met Asp Leu Arg Lys Gln Tyr Gly Asp Ala Phe Glu Lys Arg
 215 220 225
 cac ggt gtg cgt ctg ggc ttt atg tct ttc tac atc aag gcc gtg gtc 3879
 His Gly Val Arg Leu Gly Phe Met Ser Phe Tyr Ile Lys Ala Val Val
 230 235 240 245
 gaa gcg ctg aag cgt tat cca gaa gtc aac gcc tct atc gat ggc gaa 3927
 Glu Ala Leu Lys Arg Tyr Pro Glu Val Asn Ala Ser Ile Asp Gly Glu
 250 255 260
 gac gtg gtg tac cac aac tat ttc gat gtg agt att gcc gtc tct acg 3975
 Asp Val Val Tyr His Asn Tyr Phe Asp Val Ser Ile Ala Val Ser Thr
 265 270 275
 cca cgc gga ctg gtg acg cct gtc ctg cgt gac gtt gat gcg ctg agc 4023
 Pro Arg Gly Leu Val Thr Pro Val Leu Arg Asp Val Asp Ala Leu Ser
 280 285 290
 atg gct gac atc gag aag aaa att aaa gaa ctg gca gtg aaa ggc cgt 4071
 Met Ala Asp Ile Glu Lys Lys Ile Lys Glu Leu Ala Val Lys Gly Arg
 295 300 305
 gac ggc aag ctg acg gtt gac gat ctg acg ggc ggt aac ttt acc atc 4119
 Asp Gly Lys Leu Thr Val Asp Asp Leu Thr Gly Gly Asn Phe Thr Ile
 310 315 320 325
 acc aac ggt ggt gtg ttc ggt tgc ctg atg tct acg cca atc atc aac 4167
 Thr Asn Gly Gly Val Phe Gly Ser Leu Met Ser Thr Pro Ile Ile Asn
 330 335 340
 ccg cca cag agc gcg att ctg ggc atg cac gcc att aaa gat cgt cct 4215
 Pro Pro Gln Ser Ala Ile Leu Gly Met His Ala Ile Lys Asp Arg Pro
 345 350 355
 atg gcg gtc aat ggt cag gtt gtg atc ctg cca atg atg tac ctg gct 4263
 Met Ala Val Asn Gly Gln Val Val Ile Leu Pro Met Met Tyr Leu Ala
 360 365 370
 ctc tcc tac gat cac cgt tta atc gat ggt cgt gaa tct gtc ggc tat 4311
 Leu Ser Tyr Asp His Arg Leu Ile Asp Gly Arg Glu Ser Val Gly Tyr
 375 380 385
 ctg gtc gcg gtg aaa gag atg ctg gaa gat ccg gcg cgt ctg ctg ctg 4359
 Leu Val Ala Val Lys Glu Met Leu Glu Asp Pro Ala Arg Leu Leu Leu
 390 395 400 405
 gat gtc tgattcatca ctgggcacgc gttgcgtgcc caatctcaat actcttttca 4415
 Asp Val
 gatctgaatg gatagaacat c atg aac tta cac gaa tac cag gct aaa cag 4466
 Met Asn Leu His Glu Tyr Gln Ala Lys Gln
 1 5 10
 ctg ttt gca cgg tat ggc atg cca gca ccg acc ggc tac gcc tgt act 4514
 Leu Phe Ala Arg Tyr Gly Met Pro Ala Pro Thr Gly Tyr Ala Cys Thr
 15 20 25
 aca cca cgt gaa gca gaa gaa gcg gca tgc aaa atc ggt gca 4556
 Thr Pro Arg Glu Ala Glu Glu Ala Ala Ser Lys Ile Gly Ala
 30 35 40

<210> 2

<211> 39

<212> PRT

<213> Enterobacter agglomerans

<400> 2

Ala Phe Ser Val Phe Arg Cys His Ser Ile Met Asn Cys Val Ser Val
 1 5 10 15
 Cys Pro Lys Gly Leu Asn Pro Thr Arg Ala Ile Gly His Ile Lys Ser
 20 25 30
 Met Leu Leu Gln Arg Ser Ala
 35

<210> 3

<211> 935

<212> PRT

<213> Enterobacter agglomerans

<400> 3

Met Gln Asn Ser Ala Met Lys Pro Trp Leu Asp Ser Ser Trp Leu Ala
 1 5 10 15
 Gly Ala Asn Gln Ser Tyr Ile Glu Gln Leu Tyr Glu Asp Phe Leu Thr
 20 25 30
 Asp Pro Asp Ser Val Asp Ala Val Trp Arg Ser Met Phe Gln Gln Leu
 35 40 45
 Pro Gly Thr Gly Val Lys Pro Glu Gln Phe His Ser Ala Thr Arg Glu
 50 55 60
 Tyr Phe Arg Arg Leu Ala Lys Asp Ala Ser Arg Tyr Thr Ser Ser Val
 65 70 75 80
 Thr Asp Pro Ala Thr Asn Ser Lys Gln Val Lys Val Leu Gln Leu Ile
 85 90 95
 Asn Ala Phe Arg Phe Arg Gly His Gln Glu Ala Asn Leu Asp Pro Leu
 100 105 110
 Gly Leu Trp Lys Gln Asp Arg Val Ala Asp Leu Asp Pro Ala Phe His
 115 120 125
 Asp Leu Thr Asp Ala Asp Phe Gln Glu Ser Phe Asn Val Gly Ser Phe
 130 135 140
 Ala Ile Gly Lys Glu Thr Met Lys Leu Ala Asp Leu Phe Asp Ala Leu
 145 150 155 160
 Lys Gln Thr Tyr Cys Gly Ser Ile Gly Ala Glu Tyr Met His Ile Asn
 165 170 175
 Asn Thr Glu Glu Lys Arg Trp Ile Gln Gln Arg Ile Glu Ser Gly Ala
 180 185 190
 Ser Gln Thr Ser Phe Ser Gly Glu Glu Lys Lys Gly Phe Leu Lys Glu
 195 200 205
 Leu Thr Ala Ala Glu Gly Leu Glu Lys Tyr Leu Gly Ala Lys Phe Pro
 210 215 220
 Gly Ala Lys Arg Phe Ser Leu Glu Gly Gly Asp Ala Leu Val Pro Met
 225 230 235 240
 Leu Arg Glu Met Ile Arg His Ala Gly Lys Ser Gly Thr Arg Glu Val
 245 250 255
 Val Leu Gly Met Ala His Arg Gly Arg Leu Asn Val Leu Ile Asn Val
 260 265 270

Leu Gly Lys Lys Pro Gln Asp Leu Phe Asp Glu Phe Ser Gly Lys His
 275 280 285
 Lys Glu His Leu Gly Thr Gly Asp Val Lys Tyr His Met Gly Phe Ser
 290 295 300
 Ser Asp Ile Glu Thr Glu Gly Leu Val His Leu Ala Leu Ala Phe
 305 310 315 320
 Asn Pro Ser His Leu Glu Ile Val Ser Pro Val Val Met Gly Ser Val
 325 330 335
 Arg Ala Arg Leu Asp Arg Leu Ala Glu Pro Val Ser Asn Lys Val Leu
 340 345 350
 Pro Ile Thr Ile His Gly Asp Ala Ala Val Ile Gly Gln Gly Val Val
 355 360 365
 Gln Glu Thr Leu Asn Met Ser Gln Ala Arg Gly Tyr Glu Val Gly Gly
 370 375 380
 Thr Val Arg Ile Val Ile Asn Asn Gln Val Gly Phe Thr Thr Ser Asn
 385 390 395 400
 Pro Lys Asp Ala Arg Ser Thr Pro Tyr Cys Thr Asp Ile Gly Lys Met
 405 410 415
 Val Leu Ala Pro Ile Phe His Val Asn Ala Asp Asp Pro Glu Ala Val
 420 425 430
 Ala Phe Val Thr Arg Leu Ala Leu Asp Tyr Arg Asn Thr Phe Lys Arg
 435 440 445
 Asp Val Phe Ile Asp Leu Val Cys Tyr Arg Arg His Gly His Asn Glu
 450 455 460
 Ala Asp Glu Pro Ser Ala Thr Gln Pro Leu Met Tyr Gln Lys Ile Lys
 465 470 475 480
 Lys His Pro Thr Pro Arg Lys Ile Tyr Ala Asp Arg Leu Glu Gly Glu
 485 490 495
 Gly Val Ala Ser Gln Glu Asp Ala Thr Glu Met Val Asn Leu Tyr Arg
 500 505 510
 Asp Ala Leu Asp Ala Gly Glu Cys Val Val Pro Glu Trp Arg Pro Met
 515 520 525
 Ser Leu His Ser Phe Thr Trp Ser Pro Tyr Leu Asn His Glu Trp Asp
 530 535 540
 Glu Pro Tyr Pro Ala Gln Val Asp Met Lys Arg Leu Lys Glu Leu Ala
 545 550 555 560
 Leu Arg Ile Ser Gln Val Pro Glu Gln Ile Glu Val Gln Ser Arg Val
 565 570 575
 Ala Lys Ile Tyr Asn Asp Arg Lys Leu Met Ala Glu Gly Glu Lys Ala
 580 585 590
 Phe Asp Trp Gly Gly Ala Glu Asn Leu Ala Tyr Ala Thr Leu Val Asp
 595 600 605
 Glu Gly Ile Pro Val Arg Leu Ser Gly Glu Asp Ser Gly Arg Gly Thr
 610 615 620
 Phe Phe His Arg His Ala Val Val His Asn Gln Ala Asn Gly Ser Thr
 625 630 635 640
 Tyr Thr Pro Leu His His Ile His Asn Ser Gln Gly Glu Phe Lys Val
 645 650 655
 Trp Asp Ser Val Leu Ser Glu Glu Ala Val Leu Ala Phe Glu Tyr Gly
 660 665 670
 Tyr Ala Thr Ala Glu Pro Arg Val Leu Thr Ile Trp Glu Ala Gln Phe
 675 680 685
 Gly Asp Phe Ala Asn Gly Ala Gln Val Val Ile Asp Gln Phe Ile Ser
 690 695 700

Ser Gly Glu Gln Lys Trp Gly Arg Met Cys Gly Leu Val Met Leu Leu
 705 710 715 720
 Pro His Gly Tyr Glu Gly Gln Gly Pro Glu His Ser Ser Ala Arg Leu
 725 730 735
 Glu Arg Tyr Leu Gln Leu Cys Ala Glu Gln Asn Met Gln Val Cys Val
 740 745 750
 Pro Ser Thr Pro Ala Gln Val Tyr His Met Leu Arg Arg Gln Ala Leu
 755 760 765
 Arg Gly Met Arg Arg Pro Leu Val Val Met Ser Pro Lys Ser Leu Leu
 770 775 780
 Arg His Pro Leu Ala Ile Ser Ser Leu Asp Glu Leu Ala Asn Gly Ser
 785 790 795 800
 Phe Gln Pro Ala Ile Gly Glu Ile Asp Asp Leu Asp Pro Gln Gly Val
 805 810 815
 Lys Arg Val Val Leu Cys Ser Gly Lys Val Tyr Tyr Asp Leu Leu Glu
 820 825 830
 Gln Arg Arg Lys Asp Glu Lys Thr Asp Val Ala Ile Val Arg Ile Glu
 835 840 845
 Gln Leu Tyr Pro Phe Pro His Gln Ala Val Gln Glu Ala Leu Lys Ala
 850 855 860
 Tyr Ser His Val Gln Asp Phe Val Trp Cys Gln Glu Glu Pro Leu Asn
 865 870 875 880
 Gln Gly Ala Trp Tyr Cys Ser Gln His His Phe Arg Asp Val Val Pro
 885 890 895
 Phe Gly Ala Thr Leu Arg Tyr Ala Gly Arg Pro Ala Ser Ala Ser Pro
 900 905 910
 Ala Val Gly Tyr Met Ser Val His Gln Gln Gln Gln Asp Leu Val
 915 920 925
 Asn Asp Ala Leu Asn Val Asn
 930 935

<210> 4

<211> 407

<212> PRT

<213> Enterobacter agglomerans

<400> 4

Met Ser Ser Val Asp Ile Leu Val Pro Asp Leu Pro Glu Ser Val Ala
 1 5 10 15
 Asp Ala Thr Val Ala Thr Trp His Lys Lys Pro Gly Asp Ala Val Ser
 20 25 30
 Arg Asp Glu Val Ile Val Glu Ile Glu Thr Asp Lys Val Val Leu Glu
 35 40 45
 Val Pro Ala Ser Ala Asp Gly Val Leu Glu Ala Val Leu Glu Asp Glu
 50 55 60
 Gly Ala Thr Val Thr Ser Arg Gln Ile Leu Gly Arg Leu Lys Glu Gly
 65 70 75 80
 Asn Ser Ala Gly Lys Glu Ser Ser Ala Lys Ala Glu Ser Asn Asp Thr
 85 90 95
 Thr Pro Ala Gln Arg Gln Thr Ala Ser Leu Glu Glu Glu Ser Ser Asp
 100 105 110
 Ala Leu Ser Pro Ala Ile Arg Arg Leu Ile Ala Glu His Asn Leu Asp
 115 120 125
 Ala Ala Gln Ile Lys Gly Thr Gly Val Gly Gly Arg Leu Thr Arg Glu

130 135 140
 Asp Val Glu Lys His Leu Ala Asn Lys Pro Gln Ala Glu Lys Ala Ala
 145 150 155 160
 Ala Pro Ala Ala Gly Ala Ala Thr Ala Gln Gln Pro Val Ala Asn Arg
 165 170 175
 Ser Glu Lys Arg Val Pro Met Thr Arg Leu Arg Lys Arg Val Ala Glu
 180 185 190
 Arg Leu Leu Glu Ala Lys Asn Ser Thr Ala Met Leu Thr Thr Phe Asn
 195 200 205
 Glu Ile Asn Met Lys Pro Ile Met Asp Leu Arg Lys Gln Tyr Gly Asp
 210 215 220
 Ala Phe Glu Lys Arg His Gly Val Arg Leu Gly Phe Met Ser Phe Tyr
 225 230 235 240
 Ile Lys Ala Val Val Glu Ala Leu Lys Arg Tyr Pro Glu Val Asn Ala
 245 250 255
 Ser Ile Asp Gly Glu Asp Val Val Tyr His Asn Tyr Phe Asp Val Ser
 260 265 270
 Ile Ala Val Ser Thr Pro Arg Gly Leu Val Thr Pro Val Leu Arg Asp
 275 280 285
 Val Asp Ala Leu Ser Met Ala Asp Ile Glu Lys Lys Ile Lys Glu Leu
 290 295 300
 Ala Val Lys Gly Arg Asp Gly Lys Leu Thr Val Asp Asp Leu Thr Gly
 305 310 315 320
 Gly Asn Phe Thr Ile Thr Asn Gly Gly Val Phe Gly Ser Leu Met Ser
 325 330 335
 Thr Pro Ile Ile Asn Pro Pro Gln Ser Ala Ile Leu Gly Met His Ala
 340 345 350
 Ile Lys Asp Arg Pro Met Ala Val Asn Gly Gln Val Val Ile Leu Pro
 355 360 365
 Met Met Tyr Leu Ala Leu Ser Tyr Asp His Arg Leu Ile Asp Gly Arg
 370 375 380
 Glu Ser Val Gly Tyr Leu Val Ala Val Lys Glu Met Leu Glu Asp Pro
 385 390 395 400
 Ala Arg Leu Leu Leu Asp Val
 405

<210> 5
 <211> 40
 <212> PRT
 <213> Enterobacter agglomerans

<400> 5
 Met Asn Leu His Glu Tyr Gln Ala Lys Gln Leu Phe Ala Arg Tyr Gly
 1 5 10 15
 Met Pro Ala Pro Thr Gly Tyr Ala Cys Thr Thr Pro Arg Glu Ala Glu
 20 25 30
 Glu Ala Ala Ser Lys Ile Gly Ala
 35 40

<210> 6
 <211> 30
 <212> DNA
 <213> Artificial Sequence

~~<223>~~ Description of Artificial Sequence: primer

gtcgacaata/gccygaatct gttctggtcg

<210> 7

<212> DNA

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer

aagcttatcg acgctcccct ccccaccgtt

30

[illegible]